

## EDITORIAL

**MAST CELLS CONTRIBUTE TO CORONAVIRUS-INDUCED INFLAMMATION:  
NEW ANTI-INFLAMMATORY STRATEGY**S.K. KRITAS<sup>1</sup>, G. RONCONI<sup>2</sup>, AL. CARAFFA<sup>3</sup>, C.E. GALLENDA<sup>4</sup>, R. ROSS<sup>5</sup> and P. CONTI<sup>6</sup>

<sup>1</sup>Department of Microbiology and Infectious Diseases, School of Veterinary Medicine, Aristotle University of Thessaloniki, Macedonia, Greece; <sup>2</sup>Clinica dei Pazienti del Territorio, Fondazione Policlinico Gemelli, Rome, Italy; <sup>3</sup>School of Pharmacy, University of Camerino, Camerino, Italy; <sup>4</sup>Department of Biomedical Sciences and Specialist Surgery, Section of Ophthalmology, University of Ferrara, Ferrara, Italy; <sup>5</sup>University of Pennsylvania School of Veterinary Medicine, Philadelphia, PA, USA; <sup>6</sup>Postgraduate Medical School, University of Chieti, Chieti, Italy

Received January 21, 2020 – Accepted February 4, 2020

**Coronavirus, which can cause respiratory syndrome, to date has affected over seventeen thousand individuals, especially in China. Coronavirus is interspecies and can also be transmitted from man to man, with an incubation ranging from 1 to 14 days. Human coronavirus infections can induce not only mild to severe respiratory diseases, but also inflammation, high fever, cough, acute respiratory tract infection and dysfunction of internal organs that may lead to death. Coronavirus infection (regardless of the various types of corona virus) is primarily attacked by immune cells including mast cells (MCs), which are located in the submucosa of the respiratory tract and in the nasal cavity and represent a barrier of protection against microorganisms. The virus activates MCs which release early inflammatory chemical compounds including histamine and protease; while late activation provokes the generation of pro-inflammatory IL-1 family members including IL-1 and IL-33. Here, we propose for the first time that inflammation by coronavirus may be inhibited by anti-inflammatory cytokines belonging to the IL-1 family members.**

Viral infections are certainly increasing globally; they can cause an epidemic spread, but also a pandemic one in different countries, as was the case with the avian influenza virus H5N1 (mortality 35%), as well as severe acute respiratory syndrome (SARS) (mortality 10%) (1). Middle East respiratory syndrome caused by coronavirus first appeared in 2012, and since then over seventeen thousand patients have been infected worldwide (especially in China) of whom about 900 have died (a number destined to increase), according to the World Health Organization (WHO) (2). There are several types of coronavirus,

they can be transformed and become interspecies and have an incubation ranging approximately from 1 to 14 days (1). They are of unknown origin, although may originate from various animals such as snakes, bats, pigs, or dromedary camels, which are the main host, and in these animals can cause acute diarrhea, gastroenteritis, villous atrophy, poor absorption and death (2). Human-to-human virus transmission also occurs in healthcare facilities and communities. Coronaviruses often have highly pathogenic activity in animals, but today it has been seen that, with their particular pathogenic proteins, they can cause

*Key words: coronavirus; immunity; infection; inflammation; mast cells*

*Corresponding Author:*

Prof. Pio Conti,  
Postgraduate Medical School,  
University of Chieti,  
Viale Unità d'Italia 73, Chieti, Italy  
Tel./Fax: +39 0871 574136  
e-mail: pconti@unich.it

0393-974X (2020)

Copyright © by BIOLIFE, s.a.s.

This publication and/or article is for individual use only and may not be further reproduced without written permission from the copyright holder.

Unauthorized reproduction may result in financial and other penalties  
**DISCLOSURE: ALL AUTHORS REPORT NO CONFLICTS OF  
INTEREST RELEVANT TO THIS ARTICLE.**



induced type 1 Treg cells produce anti-inflammatory IL-10, protecting the body from the most aggressive forms of infection. The development of therapeutic anti-viral agents based on strengthening the immune system has been one of the main objectives of modern medicine. The immune response to the virus can be enhanced with the administration of antigens, adjuvants and vaccines. The vaccines that normally protect us from bacterial and viral infections, in the case of coronavirus, are not currently available and the experimental ones have not proved to be effective, although some vaccines have exerted a slight protective effect (13). We believe that coronavirus infection, as well as other pathogenic microorganism infections, can be combatted by preventing contagion and boosting the immune system.

#### *Mast cells*

Mast cells (MCs) are immune cells derived from

hematopoietic precursor cells (bone marrow CD34<sup>+</sup>), which mature and reside in virtually all vascularized tissue (14). MCs are located perivascularly in proximity to neurons, and their differentiation and proliferation are regulated by the stem cell factor (SCF), which binds the surface kit-receptor. After activation, MCs produce inflammatory mediators, including proteases and pro-inflammatory cytokines, participating in host response, for example, in skin diseases, and asthma (14). MCs are involved in the innate and adaptive immune systems, playing a role in autoimmunity, infections, tissue damage, and inflammatory signals (15). MCs express TLR which can be quickly recognized by pathogens, including viruses that activate them in the airways and respiratory tract, causing serious damage. When properly activated, MCs generate biologically active substances including chemokines and cytokines without degranulation. MCs are responsible for

**Fig. 2.** *Coronavirus binding Toll-like receptor; activating mast cell which secretes IL-1 family members (IL-1, IL-33), IL-6 and TNF. These cytokines along with coronavirus activate macrophages. This stimulation provokes dual effects: one being airway inflammation and fever, and the other production of IL-37 which inhibits airway inflammation and fever.*

allergic reactions, but also participate in inflammation and defend the body against bacterial helminthic and viral infections. MCs can be classically activated by IgE and specific antigen, but also by bacteria and viruses. IgE stimulates MCs via FcεRI high-affinity receptor ( $1 \times 10^{10}M^{-1}$ ), which bind to the FcεRI and cause receptor aggregation and a series of biochemical events (14). Activated MCs immediately release chemical mediators and/or late production of pro-inflammatory proteins such as cytokines and chemokines (16). Therefore, virus-activated MCs may provoke the release, after seconds, of stored chemical mediators such as histamine, tryptases and chymase. In addition, after hours of incubation, activated MCs secrete synthesized inflammatory cytokines including IL-6, IL-1, IL-31, IL-33 and TNF, and chemokines CC5, CCL2, MCP-1 and CXCL8 which attract white blood cells to the inflammatory sites (16). All these compounds have a major role in inflammation. MCs produce, store and synthesize highly inflammatory TNF, both intra-cellular and extra-cellular. MC innate immune cells are real sentinels of the human body, ready to immediately attack the external invaders which can be harmful to cells, tissues and organs. MCs release IL-33, a member of the IL-1 family of cytokines, which is considered a cytokine called "alarmin". MCs are peripherally located near dendritic cells, in communication with sensory nerves that cross-talk with brain cells, including microglia. MCs protect the body from infections caused by bacterium and viruses (17). These cells, communicating with the external environment, neutralize various pathogens, but in doing so they also produce pro-inflammatory compounds which can be harmful for the body. Stressful and chronic conditions predispose humans to pathogenic infections including those caused by coronavirus. In fact, neurotransmitters, including substance P, that are generated in stress, can activate immune cells such as MCs to produce pro-inflammatory cytokines and chemical mediators of inflammation (18).

Experiments with coronavirus in mice have shown that this pathogenic RNA microorganism can activate the cells of both innate and adaptive immune systems, including MCs. These cells are located in

the submucosa of the respiratory tract and in the nasal cavity and represent the protective cells that are at the forefront in fighting viral and bacterial agents (19). They are involved in many biological processes where they induce the secretion of a broad spectrum of cytokines and chemokines. In addition, viruses activating MCs cause the release of some specific chemokines, such as the ligand 5 (CCL5), which attracts the CD8 T immune cells that defend the lung tissue and fight viral infection (20). Epithelial cells represent a physical barrier to the entry of microorganisms including viruses. The cross-talk between viruses and epithelial cells causes the production of immune mediators in order to defend the body. Viruses develop particular mechanisms to invade the body and immune cells including MCs. The host response to RNA virus invasion activates TLR3 on MCs with an anti-viral IFN production (positive effect) (21); but often the virus only causes sensitization of MCs with the synthesis of IgE that bind to the FcεRI receptor and trigger a violent inflammatory reaction (negative effect). Therefore, the function of MCs towards viral infections is not yet clear. However, RNA viruses stimulate MCs to produce IFN gamma e CXCL8 chemokines, resulting in the recruitment of NK cells that also produce type I interferons (IFNs) which are anti-viral cytokines. In addition, type I IFNs enhance the cytotoxic activity of NK cells against virus-infected cells (22). So, in this case, MCs are anti-viral immune cells through the above mechanism. On the other hand, viruses stimulate mucosa MCs to release pro-inflammatory cytokines such as TNF, IL-1, IL-6 and proteases, which aggravate the inflammatory state.

Virus-activated MCs produce histamine, prostaglandin D<sub>2</sub> (PGD<sub>2</sub>), and leukotriene C<sub>4</sub> (LTC<sub>4</sub>) which induces acute bronchoconstriction and lung inflammation. Therefore, we can conclude that viral infections can activate MCs which respond with dual effects, a positive one helping the immune system to fight infection, and a negative one causing the release of chemical mediators of inflammation and the secretion of pro-inflammatory cytokines such as IL-1, IL-33, IL-18, and TNF, aggravating the pathological state of the patient.

IL-37 is a member of the IL-1 family of cytokines,

which is activated by caspase-1 and suppresses transcription of pro-inflammatory genes and immune responses. Since IL-37 is an inhibitor of IL-1, a potent pro-inflammatory cytokine (23), we believe that this anti-inflammatory cytokine may suppress fever and inflammation provoked by coronavirus, so as to reduce the number of deaths.

## REFERENCES

- Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med* 2020; doi: 10.1056/NEJMoa2001316. [Epub ahead of print].
- Holshue ML, DeBolt C, Lindquist S, et al. First case of 2019 novel coronavirus in the United States. *N Engl J Med* 2020; doi: 10.1056/NEJMoa2001191. [Epub ahead of print].
- Conti P. Sistema immunitario forte contro il virus. *Il Centro* 2020; Gennaio 31:p.5.
- Phan LT, Nguyen TV, Luong QC, et al. Importation and human-to-human transmission of a novel coronavirus in Vietnam. *N Engl J Med* 2020; doi: 10.1056/NEJMc2001272. [Epub ahead of print].
- Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020; doi: 10.1056/NEJMoa2001017 [Epub ahead of print].
- Shokri S, Mahmoudvand S, Taherkhani R, Farshadpour F. Modulation of the immune response by Middle East respiratory syndrome coronavirus. *J Cell Physiol* 2019; 234(3):2143-51.
- Morfopoulou S, Brown JR, Davies EG, et al. human coronavirus oc43 associated with fatal encephalitis. *N Engl J Med* 2016; 375(5):497-8. doi: 10.1056/NEJMc1509458.
- Perlman S. Another decade, another coronavirus. *N Engl J Med* 2020; doi: 10.1056/NEJMe2001126. [Epub ahead of print].
- Oh MD, Park WB, Choe PG, et al. Viral load kinetics of MERS coronavirus infection. *N Engl J Med* 2016; 375(13):1303-5. doi: 10.1056/NEJMc1511695.
- Li G, Fan Y, Lai Y, et al. Coronavirus infections and immune responses. *J Med Virol* 2020; doi: 10.1002/jmv.25685. [Epub ahead of print] Review.
- Doki T, Tarusawa T, Hohdatsu T, Takano T. In vivo antiviral effects of U18666A against Type I feline infectious peritonitis virus. *Pathogens* 2020; 9(1). doi: 10.3390/pathogens9010067.
- de Aquino MT, Puntambekar SS, Savarin C, et al. Role of CD25(+) CD4(+) T cells in acute and persistent coronavirus infection of the central nervous system. *Virology* 2013; 447(1-2):112-20.
- Enjuanes L, Zuñiga S, Castaño-Rodríguez C, et al. molecular basis of coronavirus virulence and vaccine development. *Adv Virus Res* 2016; 96:245-86.
- Galli SJ, Kalesnikoff J, Grimbaldston MA, et al. Mast cells as “tunable” effector and immunoregulatory cells: recent advances. *Annu Rev Immunol* 2005; 23:749-86. Review.
- Taracanova A, Tsilioni I, Conti P, Norwitz ER, Leeman SE, Theoharides TC. Substance P and IL-33 administered together stimulate a marked secretion of IL-1 $\beta$  from human mast cells, inhibited by methoxyluteolin. *Proc Natl Acad Sci U S A* 2018; 115(40):E9381-E9390.
- Gallenga CE, Pandolfi F, Caraffa AI, et al. Interleukin-1 family cytokines and mast cells: activation and inhibition. *J Biol Regul Homeost Agents* 2019; 33(1):1-6.
- Antonopoulos D, Tsilioni I, Balatsos NAA, Gourgoulisian KI, Theoharides TC. The mast cell - neurofibromatosis connection. *J Biol Regul Homeost Agents* 2019; 33(3):657-59.
- Conti P, Lauritano D, Caraffa A, et al. New insight into systemic mastocytosis mediated by cytokines IL-1 $\beta$  and IL-33: Potential inhibitory effect of IL-37. *Eur J Pharmacol* 2019; 858:172473.
- Marshall JS, Portales-Cervantes L, Leong E. Mast cell responses to viruses and pathogen products. *Int J Mol Sci* 2019; 20(17). pii: E4241.
- Podlech J, Ebert S, Becker M, Reddehase MJ, Stassen M, Lemmermann NA. Mast cells: innate attractors recruiting protective CD8 T cells to sites of cytomegalovirus infection. *Med Microbiol Immunol* 2015; 204(3):327-34.
- Meng D, Huo C, Wang M, et al. Influenza A viruses replicate productively in mouse mastocytoma cells (P815) and trigger pro-inflammatory cytokine and chemokine production through TLR3 signaling pathway. *Front Microbiol* 2017; 7:2130.
- Oki S, Miyake S. Invariant natural killer T (iNKT)

cells in asthma: a novel insight into the pathogenesis of asthma and the therapeutic implication of glycolipid ligands for allergic diseases. *Allergol Int* 2007; 56(1):7-14.

23. Cavalli G, Dinarello CA. Suppression of inflammation and acquired immunity by IL-37. *Immunol Rev* 2018; 281(1):179-90.

PROOF